

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Acibenzolar-S-methyl may prevent vector-mediated flavescence dorée phytoplasma transmission, but is ineffective in inducing recovery of infected grapevines**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1637742> since 2017-07-17T12:26:33Z

*Published version:*

DOI:10.1002/ps.4303

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Miliordos D.E, Galetto L, Ferrari E, Pegoraro M, Marzachì C, Bosco, D..  
Acibenzolar-S-methyl may prevent vector-mediated flavescence dorée  
phytoplasma transmission, but is ineffective in inducing recovery of infected  
grapevines. PEST MANAGEMENT SCIENCE. 73 pp: 534-540.  
DOI: 10.1002/ps.4303

The publisher's version is available at:

<http://onlinelibrary.wiley.com/doi/10.1002/ps.4303/fullpdf>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/1637742>



**ACIBENZOLAR-S-METHYL MAY PREVENT VECTOR-MEDIATED  
FLAVESCENCE DORÉE PHYTOPLASMA TRANSMISSION, BUT  
IS INEFFECTIVE IN INDUCING RECOVERY OF INFECTED  
GRAPEVINES**



Journal:	<i>Pest Management Science</i>
Manuscript ID	PM-16-0039.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Miliordos, Dimitrios; Universita di Torino, Dipartimento di Scienze Agrarie, Forestali e Alimentari Galletto, Luciana; Consiglio Nazionale delle Ricerche, Istituto di Protezione Sostenibile delle Piante Ferrari, Ester; Universita di Torino, Dipartimento di Scienze Agrarie, Forestali e Alimentari Pegoraro, Mattia; Universita di Torino, Dipartimento di Scienze Agrarie, Forestali e Alimentari Marzachi, Cristina; Consiglio Nazionale delle Ricerche, Istituto di Protezione Sostenibile delle Piante Bosco, Domenico; Universita di Torino, Dipartimento di Scienze Agrarie, Forestali e Alimentari; Consiglio Nazionale delle Ricerche, Istituto di Protezione Sostenibile delle Piante
Key Words:	Resistance elicitor, Flavescence dorée, Scaphoideus titanus, grapevine, phytoplasma transmission, recovery

SCHOLARONE™  
Manuscripts

**ACIBENZOLAR-S-METHYL MAY PREVENT VECTOR-MEDIATED FLAVESCENCE  
DORÉE PHYTOPLASMA TRANSMISSION, BUT IS INEFFECTIVE IN INDUCING  
RECOVERY OF INFECTED GRAPEVINES**

Dimitrios E Miliordos<sup>a</sup>, Luciana Galetto<sup>b</sup>, Ester Ferrari<sup>a</sup>, Mattia Pegoraro<sup>a</sup>, Cristina Marzachi<sup>b</sup>,  
Domenico Bosco<sup>a,b\*</sup>

<sup>a</sup>DISAFA – Entomologia, Università degli Studi di Torino, Largo Paolo Braccini 2, 10095 Grugliasco (TO), Italy

<sup>b</sup>CNR – Istituto di Protezione Sostenibile delle Piante, IPSP, Strada delle Cacce 73, 10135 (TO), Italy

\*Correspondence to : Domenico Bosco, DISAFA – Entomologia, Università degli Studi di Torino, Largo Paolo Braccini 2, 10095 Grugliasco (TO), Italy. Email: domenico.bosco@unito.it

**Abstract**

**BACKGROUND:** Acibenzolar-S-methyl (BTH), a functional analog of Salicylic Acid (SA), is known to elicit a systemic resistance across a broad range of plant–pathogen interactions but, so far, it was not tested against Flavescence dorée (FDP), one of the most devastating grapevine diseases. Aims of this work were to evaluate the activity of BTH in preventing FDP transmission by the insect vector and in inducing recovery of infected grapevines.

**RESULTS:** Repeated applications of 2 mM BTH to test grapevine cuttings (cv Barbera) exposed to adults of the infectious vector, *Scaphoideus titanus* Ball, reduced the rate of infected plants. The effect was not recorded following similar BTH applications to highly susceptible young *in vitro* propagated vines. A high natural recovery rate (more than 70%) was observed over a three-year-period in field-infected grapevines of the same cv. Under these conditions, BTH repeated applications over the whole period clearly failed to increase recovery of field-infected grapevines.

**CONCLUSION:** Following a three-year experiment, it can be concluded that, although high doses and repeated applications of BTH reduced vector transmission of FDP, BTH was ineffective in inducing recovery of FDP-infected grapevines cv Barbera under field conditions.

**Key words:** Resistance elicitor, Flavescence dorée, *Scaphoideus titanus*, grapevine, transmission, recovery

## 1 INTRODUCTION

The use of induced resistance to increase the effectiveness of plant natural defences is a promising environment-friendly tool for plant protection. Its introduction into agricultural practices can reduce the use of chemicals for pathogen control, thus contributing to the development of sustainable agriculture. The phenomenon of induced resistance has been described as Systemic Acquired Resistance (SAR) and as Induced Systemic Resistance (ISR). SAR is a long-lasting kind of induced resistance effective against a broad spectrum of pathogens<sup>1-3</sup>. In the vast majority of cases, it depends on the early increase of the endogenously synthesized signal salicylic acid (SA)<sup>4</sup> and is correlated with the immediate expression of a specific set of genes called “SAR genes”, which includes genes coding for pathogenesis-related (PR) proteins<sup>5-7</sup>. ISR is controlled by a signaling pathway in which the phytohormones jasmonic acid (JA) and ethylene play key roles as revealed by the use of SA, JA and ethylene signaling mutant plants<sup>8</sup>. ISR is a resistance active against various bacterial, viral, fungal and nematode pathogens and has been found active in many plant species.

The use of SAR and ISR to cope with crop diseases and improve integrated pest management seems promising. BTH and INA (2,6-dichloroisonicotinic acid) are by far the best studied chemical elicitors available. Both are considered functional analogs of SA, and elicit a systemic form of induced resistance across a broad range of plant–pathogen interactions<sup>9-10</sup>. The efficacy of BTH as a resistance inducer has been demonstrated in many crops to a wide variety of pathogens also under field conditions. Its effect lasts longer in monocots, such as wheat, than in dicots crops in

1 which repeated applications are required to extend protection over time<sup>11-12</sup>. Indeed, BTH was  
2 originally marketed as a tool to control powdery mildew of wheat and barley in Europe<sup>11</sup>.

3       Despite the almost 30 year long BTH application under different field conditions, studies of  
4 systemic resistance induced by BTH against phytoplasmas are in their infancy. So far, most BTH  
5 applications have been tested on phytoplasma diseases of herbaceous plants. For instance, BTH  
6 showed a strong protection activity against X-disease phytoplasma when applied to *Arabidopsis*  
7 *thaliana* plants<sup>13</sup>. Moreover, these authors reported a reduced survival of the leafhopper vector  
8 *Colladonus montanus* (Van Duzee), when it fed on 4.8 mM BTH treated plants, suggesting that SAR  
9 in this plant may have some detrimental effect on the leafhopper *C. montanus*. BTH application to  
10 daisy plants delayed symptom development and phytoplasma ('*Candidatus* Phytoplasma asteris',  
11 strain CYP) multiplication in treated plants compared with the control ones<sup>14</sup>. As for the application  
12 of BTH to phytoplasma-infected grapevines, field treatment with resistance inducers for the  
13 control of Bois Noir phytoplasma provided encouraging results<sup>15</sup>. Flavescence dorée is a  
14 phytoplasma-associated disease of grapevine that causes severe damages in large viticultural areas  
15 of Europe, namely in Northern Italy and Southern France. In some areas, e.g. North-Western Italy,  
16 it is the main constraint to viticulture. Flavescence dorée phytoplasmas, FDP, belong to the 16Sr V-  
17 C and D ribosomal groups and are transmitted by the leafhopper *Scaphoideus titanus* Ball  
18 (Hemiptera, Cicadellidae). FDP is a quarantine organism, its control is compulsory by law, and it  
19 relies mainly on insecticide applications against the vector and rouging of infected plants. In spite  
20 of all the efforts, a satisfactory control of the disease is difficult to achieve because many disease  
21 and vector reservoirs occur in the uncultivated areas and woodlands nearby vineyards<sup>16</sup>.

22       Aims of this study were to investigate the activity of BTH in preventing Flavescence dorée  
23 phytoplasma (FDP) transmission by the vector *Scaphoideus titanus* Ball and in inducing recovery of  
24 FDP-infected grapevines. For the first goal, healthy grapevine cuttings were sprayed with BTH and

then exposed to FDP-infectious *S. titanus*. For the second goal, FDP-infected grapevines were identified in the vineyard, BTH was applied to a group of infected plants over a three-year period and recovery from FDP was recorded on BTH-treated and control grapevines.

## 2 MATERIALS AND METHODS

### 2.1 Vector and FD phytoplasma

Two species of FDP leafhopper vectors were used in the experiments: *S. titanus*, the natural vector<sup>17</sup>, and *Euscelidius variegatus* Kirschbaum, a laboratory vector, routinely used to maintain FDP under laboratory conditions<sup>18</sup>. *S. titanus*, which has one generation per year, was used to inoculate test grapevines, while *E. variegatus*, which can be continuously reared, was used to maintain under laboratory conditions the FDP-C strain<sup>19</sup> on broad bean (*Vicia faba* L.). To obtain healthy *S. titanus* nymphs, two year-old branches bearing leafhopper eggs were collected in infested vineyards during the winter period, cut into 20-30 cm long pieces and kept in plastic bags in a cold room at 4°C. To allow egg hatching, branches were caged inside insect-proof screen houses (100 x 100 x 100 cm) in a glasshouse with natural light and temperature ranging from 20 to 30 °C. Potted grapevine cuttings and healthy broad bean plants from seed were introduced in the screen house to feed the newly hatched nymphs. Insect nymphs and adults were periodically collected and used for the experiments. Healthy colonies of *E. variegatus* were reared on potted oat plants inside plexiglas and nylon cages in growth chambers at 23 °C (photoperiod L16:D8 light:dark) and periodically checked to be phytoplasma-free by nested PCR with the universal primers P1/P7<sup>20-21</sup>, followed by F1/R1(V) primers<sup>22</sup>.

FDP-C strain<sup>19</sup> was originally isolated from infected grapevines in Piemonte Region and transmitted to broad bean plants by *S. titanus*. Once the FDP was transmitted to broad beans, the

phytoplasma was maintained by insect transmission, from broad bean to broad bean, under laboratory conditions. Third and fourth instar nymph of *E. variegatus* were introduced in a plexiglas and nylon cage and fed on FDP-infected broad beans for 7 day acquisition access periods (AAP). Then, leafhoppers were transferred onto healthy oat plants for three weeks, to complete the latent period (LP), and finally transferred onto healthy broad beans for successive inoculation access periods (IAP). Inoculated broad beans were replaced twice a week and, after the IAP, they were maintained in a climatic chamber at 23-25 °C (photoperiod L16:D8 light: dark) for at least one month and observed for specific symptom development. Following the IAP, broad beans were treated with the insecticide Dichlorvos (Dedevap, Bayer Crop Science) and later served as inoculum source for healthy *S. titanus*. To obtain FDP-infectious *S. titanus*, nymphs were introduced in a plexiglas and nylon cage with four FDP-infected broad beans for a 7 day AAP. The infected broad beans were then replaced with healthy broad bean plants for 21 day LP, when infected insects were collected for grapevine inoculations.

2.2 Protection from FDP transmission by *Scaphoideus titanus* - grafted cuttings

Two year-old plants of *Vitis vinifera* cv Barbera grafted onto Kober 5BB rootstock were used for this experiment. Plants were grown in a screen house under field conditions in 10 L pots containing peat and pumice mixture (2:1). Inoculation by infectious *S. titanus* was performed at the beginning of July, when each plant had two shoots and each shoot had 8-12 leaves. For the experiment, four screen houses (two treated and two not treated) containing 20 potted grapevines were set, for a total of 80 plants. The experiment was repeated for three consecutive years: 2011, 2012 and 2013. To evaluate the resistance induced by BTH (Bion, Syngenta Crop Protection) against FDP, 40 grapevines were foliar sprayed with 2.4 mM BTH until run off point one week before the inoculation by infectious *S. titanus*. A total volume of 2 L was applied to the 40 plants and



therefore, about 50 ml of 0.5 g BTH was applied to each of the potted cuttings. The concentration of 2.4 mM was selected on the basis of the results obtained in a previous study on the activity of BTH in preventing the transmission of another phytoplasma (chrysanthemum yellows phytoplasma, '*Candidatus Phytoplasma asteris*') to daisy plants<sup>14</sup>. The application was repeated every 10 days until the end of September. Forty control grapevines were sprayed with water at the same dates.

Twenty-five FD-infectious adults obtained as described above were introduced in each of the four screen houses (on both BTH-treated and untreated grapevines), while other 20 leafhoppers were analyzed by nested PCR in order to estimate the number of FDP-positive insects. Every year, in December, all potted grapevines were pruned and transplanted in an experimental vineyard under field condition. In the following year the grapevines (BTH treated and untreated) were cultivated according to conventional practices. Moreover, BTH was applied, at the same dose, to the BTH-treated field transplanted vines, from mid-May until mid-September every 10 days. The infection status of the test plants was checked by symptom observation and PCR two months after inoculation and in the following year.

In order to exclude any effect of BTH against the leafhopper vector, two grapevines were BTH treated as described above and two were water treated. Ten newly emerged adults of *S. titanus* were caged on each vine and the survival of the leafhoppers recorded one week later.

### 2.3 Protection from FDP transmission by *Scaphoideus titanus* - *in vitro* propagated grapevines

Grapevine plants (*V. vinifera* cv. Barbera) were propagated *in vitro*<sup>23</sup> and then transplanted in 12 cm pots and grown in a greenhouse at 20-25 °C. Vines with 7-10 leaves, one month after transplant, were used for the experiment. One week before the inoculation of FDP with infectious *S. titanus*, BTH (2.4 mM) was sprayed onto leaves of 10 plants, whilst other 10 plants were water

1  
2  
3 1 treated as control. BTH was applied until the run off point, and the application was repeated every  
4  
5 2 10 days until the end of the experiment (11 weeks post inoculation). FDP-infected *S. titanus* adults  
6  
7 3 were obtained as described above and all twenty grapevines were singly exposed to four  
8  
9 4 leafhoppers for an inoculation access period (IAP) of seven days. After the inoculation period the  
10  
11 5 surviving insects were collected and stored at -20° C. Total DNA was extracted from twenty insects  
12  
13 6 and analyzed by nested PCR in order to estimate the number of the FDP-positive individuals. At  
14  
15 7 five and eleven weeks post inoculation five leaves were sampled from each test plant, DNA was  
16  
17 8 extracted and FDP was detected by PCR. Samples that tested positive by PCR were further  
18  
19 9 analyzed by q-PCR in order to estimate the phytoplasma population in BTH-treated and untreated  
20  
21 10 test grapevines.  
22  
23  
24  
25  
26  
27  
28

29 12 **2.4 Induction of recovery in field-infected grapevines**

30  
31 13 The study was carried out in a vineyard of cv Barbera at Cocconato (Asti province), Piemonte  
32  
33 14 Region, north-western Italy. The vineyard was planted in 1998 on SO4 rootstock and consisted of  
34  
35 15 76 rows for a total of about 8800 grapevines, the training system was cordon pruning and vines  
36  
37 16 were submitted to conventional agronomic practices, including fungicide and insecticide  
38  
39 17 applications. In particular, for the control of the FDP vector, insecticide was applied twice during  
40  
41 18 the vegetation season, as enforced by the law in the areas where both FDP and its vector are  
42  
43 19 present (Italian Ministry of Agriculture, 2000). The first application was done at mid-end of June,  
44  
45 20 against *S. titanus* nymphs (Actara, Syngenta Crop Protection), and the second application took  
46  
47 21 place one month later at mid or end of July against leafhopper adults with Chlorpyrifos- ethyl  
48  
49 22 (Dursban, Dow AgroSciences).  
50  
51  
52  
53  
54

55 23 FDP-infected grapevines were identified during sampling surveys at the end of June –  
56  
57 24 beginning of July in 2011. Plants with phytoplasma symptoms were sampled and diagnosed for the  
58  
59  
60

presence of FDP. According to the symptoms and the diagnostic PCR assays, 72 FDP-infected grapevines were identified for the experiment. Half of the plants, randomly distributed in the vineyard, were foliar sprayed with 2.4 mM BTH until run off point every 10 days from mid-May (mid-July in 2011) until mid-September for three consecutive years (2011, 2012 and 2013). A total volume of 15 L was applied to the 36 plants and therefore, about 400 ml of 2.4 mM BTH was applied to each plant. Control plants were sprayed with similar volumes of water. When two infected vines were next to each other, both were included in the same treatment (BTH or water), in order to avoid drifting of BTH to a neighboring control plant. Every year, at the beginning of September, each grapevine was checked for FDP symptoms and analyzed for the presence of FDP by nested PCR. In the final year, 2013, FDP positive grapevines were analyzed by quantitative real time PCR (q-PCR) in order to measure the phytoplasma load in treated and untreated test grapevines.

## 2.5 Total DNA extraction and FDP detection

Total DNA was extracted from 1.5 g of leaf veins according to a phytoplasma enrichment protocol followed by a CTAB-based extraction procedure and dissolved in 100 µl sterile double distilled water (SDW)<sup>24</sup>. Total DNA was also extracted from single leafhoppers following a procedure previously described by Marzachi *et al.*<sup>25</sup>. DNA concentration of extracts was quantified with a Nanodrop (ND 1000 Spectrophotometer; Nanodrop technologies, Inc.). Nested PCR was used for phytoplasma detection. Universal primers pair P1/P7 was used in direct PCRs<sup>20-21</sup>. Reaction products were diluted 1:40 in SDW and used as templates in nested reaction driven by primers R16(V)F1/R1 for the specific detection of 16S group V<sup>22</sup>. Amplicons were separated by electrophoresis in 1% agarose TBE (90 mM Tris- borate, 2 mM EDTA) gel and visualized under UV light after staining with ethidium bromide.

1

2

3 1

4

5 2 **2.6 Quantification of Flavescence dorée phytoplasma cells**

6

7 FDP titre in infected grapevines was measured by q-PCR as the number of FD phytoplasma

8

9

10 4 cells per ng of grapevine DNA according to the method described in Roggia *et al.*<sup>26</sup>, based on a

11

12 5 double absolute quantification assay (phytoplasma and plant DNA) designed by Marzachi and

13

14 6 Bosco<sup>27</sup>. To determine the number of FD phytoplasma cells, specific primers targeting a portion of

15

16

17 7 the *secY* gene of the FD phytoplasma were used. For the quantification of grapevine DNA, specific

18

19 8 primers targeting the 18S rDNA gene of grapevine were used. Standard curves for the absolute

20

21 9 quantification of FD phytoplasma and grapevine DNA were obtained by dilution of plasmid

22

23

24 10 p26SecYFD, containing the appropriate *secY* gene target sequence from FD-C phytoplasma, and of

25

26 11 total DNA extracted from healthy grapevine grown from seed, respectively. Dilution of grapevine

27

28

29 12 samples, reaction mix and cycling conditions were as described in the original paper<sup>26</sup>. Each

30

31 13 sample was run in triplicate in the same plate. For each PCR, water instead of DNA was used as

32

33

34 14 negative control. Threshold levels, threshold cycles and standard curves were automatically

35

36 15 calculated by the iCycler software (BioRad), v. 3.06070. Per-well baseline cycles were determined

37

38 16 automatically. Specificity of the reaction was tested by running a melting curve analysis of the

39

40 17 amplicons following each quantification reaction.

41

42

43 18

44

45 19 **2.7 Data analysis**

46

47 Chi-square (or Fisher exact test) was applied to the comparison of the proportion of

48

49

50 20 infected/recovered plants in the two treatments (BTH and water) in all the experiments. For the

51

52 21 analysis of the phytoplasma population, raw data (obtained as log) were used and the

53

54 22 concentration of FDP in each plant was expressed as the difference between the logarithm of the

55

56 23 number of FDP cells and the logarithm of ng grapevine DNA. FDP population in treated and not

57

58 24

59

60

1 treated Barbera plants was compared with t-test. For all the analyses SigmaPlot 11.0 (Systat  
2 Software, Inc., San Jose, CA) was used.

### 3 RESULTS

#### 3.1 Protection from FDP infection by *Scaphoideus titanus*

Some plants (8 among the BTH-treated and 19 among the control plants) did not survive following the transplantation into the vineyard, and therefore the results refer to the surviving plants. Only one plant tested PCR positive for FD infection the year of the inoculation. A total of 8 out of 113 (7.0 %) BTH-treated grapevines were infected by FDP after the inoculation in the three-year experiment, whilst 18 out of 101 (17.8 %) control grapevines were infected under the same conditions (Table 1). Chi-square analysis showed a significant effect of BTH in protecting from FDP infection (Chi-square = 4,803, 1 df,  $P = 0,028$ ), on pooled data collected during the three years. The analysis of the year by year results showed a significant/almost significant effect of BTH in 2012 ( $P = 0.020$ ) and in 2013 ( $P = 0.055$ ), while no effect was recorded in 2011 ( $P = 0.367$ ).

The three batches of 20 *S. titanus* specimens analyzed every year to estimate the proportion of infected insects showed 60 to 70 % of FDP-positive leafhoppers (2011, 12/20; 2012, 14/20; 2013, 12/20).

No activity of BTH against *S. titanus* was recorded, as the survival was similar on both BTH-treated and untreated vines.

#### 3.2 Protection from FD infection by *Scaphoideus titanus* – *in vitro* propagated cuttings

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1           Eight out of ten BTH-treated and six out of ten control grapevines became infected with FDP  
2 following inoculation with four leafhoppers per plant (Table 2). Thus, no differences between the  
3 treated and untreated *in vitro* propagated vines were detected (Fisher exact test,  $P = 0.628$ ).  
4 Seventeen out of 20 *S. titanus* sampled after inoculation of the test plants were FDP-positive  
5 following nested PCR assay (85%). The first FD symptoms appeared during the 5th week post  
6 inoculation, on both BTH treated and control plants.

7           Phytoplasma population in the FDP-infected grapevines, BTH-treated and untreated, was  
8 measured on leaf samples at 5 and 11 weeks post-inoculation (wpi), and at the former date FDP  
9 load was very low regardless of the treatment. At both times, an apparently higher number of FDP  
10 cells was estimated in BTH-treated plants, but these differences were not significant (t-test: 5 wpi  $P$   
11 = 0.178, 11 wpi  $P = 0.082$ ). A significant increase of FDP cells over time was recorded in both in  
12 BTH-treated (t-test,  $P = 0.004$ ) and untreated plants (t-test,  $P = 0.016$ ), demonstrating an active  
13 phytoplasma multiplication in the infected plants during the early phases of infection.

14  
15   **3.3 Induction of recovery in field-infected grapevines**

16           Two FDP-infected grapevines died during the second year of the experiment and therefore  
17 were excluded from the analysis. After the three-year field experiment, most of the FDP-infected  
18 Barbera plants underwent recovery. Fifty-eight and 73.5% of the FDP-infected grapevines were  
19 recovered (symptomless and FDP negative by PCR) at the end of the study among BTH-treated and  
20 untreated plants, respectively (Table 3). The statistical analysis showed no significant differences  
21 between the treatments (Chi-square = 1.181, 1 df,  $P = 0.277$ ) and therefore plants recovered from  
22 FDP regardless of the treatment.

23           The temporal pathways of recovery of the single vines over time are shown in  
24 Supplementary Tables 1 (BTH-treated) and 2 (control plants). Thirteen and 20 grapevines recovered

in the second year of the study in the BTH and control treatments, respectively. Nine and 7 grapevine plants recovered the third year of the study in the BTH and control treatments, respectively. Four plants, two among the BTH-treated and two among control plants, following a recovery in the second year, were newly infected in the third year (Supplementary Tables 1 and 2). The total numbers of FDP-infected grapevines during the three-years of the experiment are shown in Figure 1 (BTH-treated and control plants). In the third year, at the end of the experiment, FDP population was about 700 and 1,200 cells per ng of grapevine DNA in BTH-treated and control plants, respectively (Table 3). The difference was not statistically significant (t-test = -1,281, 19 df, P = 0.216) and thus the treatment had no effect on the phytoplasma multiplication in the infected plants.

#### 4 DISCUSSION AND CONCLUSIONS

One of the main challenges in Northern Italy and Southern France viticulture is to control/suppress the spread of FD disease. Several measures are applied, such as compulsory insecticides against the vector, rouging of infected plants, cleaning of abandoned vineyard and of wild rootstocks that can host the phytoplasma and the vector. In spite of all the efforts, the disease is still spreading in several areas where it represents the main constraint to viticulture. Recently, induction of “recovery” of the infected grapevines is being explored as a tentative solution<sup>28</sup>. Also, in the frame of integrated FD management, agronomic techniques such as severe pruning or rouging of infected plants and the application of compounds or plant symbiotic microorganisms that may act as resistance elicitors, are under evaluation<sup>29</sup>. The present study represents the first trial of BTH application on grapevines to induce resistance against FDP.

Setting of the experimental conditions (dosage and application timing) was based on previous work measuring the activity of BTH against chrysanthemum yellows phytoplasma (CYP) in

daisy plants<sup>14</sup>. However, dosage and application frequency of this experiment are unlikely to be applied in commercial vineyards; indeed, they were designed to maximize the identification of any BTH effect on FD infection.

After the three-year experiment, considering all the inoculated grafted cutting test plants, FDP-infected grapevines were significantly more numerous in the control plants than among those sprayed with BTH, thus indicating an activity of BTH in preventing transmission of FDP. However, due to the low proportion of infected plants recorded for both BTH and control experiments, and the fact that repeats of the three years (2011, 2012 and 2013) provided slightly different results, BTH activity in preventing FDP transmission to grapevines could not be proven definitively. On the other hand, most of the cuttings from *in vitro* propagated vines were successfully infected by infectious *S. titanus*, irrespective of BTH applications. The reasons for the higher susceptibility of *in vitro* derived plants compared to grafted ones may be different. Herbaceous hosts of phytoplasmas are generally very susceptible and easy to infect, as it is the case of daisy for CYP<sup>30</sup>, broad bean for FDP<sup>31</sup>, and many other herbaceous hosts for different phytoplasmas that can be routinely and efficiently transmitted by their vectors. A high susceptibility to FDP was also reported by Jarausch *et al.*<sup>32</sup> for young grapevines obtained from *in vitro* multiplication following inoculation by the vector *S. titanus*. Also, a different feeding behavior of the vector on young, tender shoots/leaves cannot be excluded and possibly explain the different transmission results. Finally, it is possible that the absence of rootstock increases susceptibility of grapevine to the pathogen. The higher infectivity of leafhoppers that inoculated *in vitro* propagated vines, besides the higher number of insects per plant, may also explain the different transmission, despite their shorter IAP. Moreover, no effect of BTH was measured on the multiplication of FDP in infected cuttings from *in vitro* propagated vines, a further indication that this molecule is not effective in eliciting resistance on these very susceptible plants.



To our knowledge, there are only two reports on the activity of BTH in preventing the transmission of phytoplasmas by the insect vectors<sup>13-14</sup>. Interestingly, the results are not consistent for the two phytoplasma-plant vector associations. While Bressan and Purcell<sup>13</sup> reported a significant effect of BTH in reducing X-disease phytoplasma transmission to *Arabidopsis thaliana* (L.) Heynh by the leafhopper *Colladonus montanus* (Van Duzee), D'Amelio *et al.*<sup>14</sup> reported no differences in transmission efficiencies of CYP to *C. carinatum* by the vector *M. quadripunctulatus*. In this work, a significant effect was found in preventing transmission of FDP to grafted grapevines and no effect in preventing FDP transmission to herbaceous *in vitro*-propagated grapevines with the vector *S. titanus*. In spite of the lack of activity in preventing CYP transmission, D'Amelio *et al.*<sup>14</sup> reported a significantly lower phytoplasma load in the BTH-treated test plants at early stages of infection, while no differences in FDP load were found between treated and untreated grapevines under our experimental conditions. The activity of BTH in preventing phytoplasma transmission is highly dependent on the phytoplasma-plant-vector association.

In the last decade, the spontaneous remission of symptoms in previously infected plants (recovery) has attracted a lot of attention for the potential consequences on economically important woody crops. In particular, recovery from phytoplasmas was reported for apple<sup>33-34</sup>, apricot<sup>35</sup>, pear<sup>36</sup> and grapevine<sup>37-39</sup>. Recovery from Bois Noir (BN), another grapevine yellows of phytoplasma origin, can be promoted by the application of resistance elicitors compounds or stress practices such as partial uprooting of the plants<sup>40-41</sup>. In the present work, repeated BTH foliar applications to FDP-infected grapevines under field conditions for three years did not increase the number of Barbera grapevines that underwent recovery compared to water-treated control plants. Indeed, a very high spontaneous recovery rate occurred in the control plants, and even more untreated grapevines underwent recovery compared to the treated ones, though the difference was not significant. Moreover, no significant difference was recorded in the number of FDP cells in

1 treated and control plants that were still infected at the end of the experiment, as measured by q-  
2 PCR. These results are in line with those showing that BTH application do not affect either the  
3 apple proliferation phytoplasma infection rate or phytoplasma multiplication in treated apple trees  
4 under controlled conditions<sup>42</sup>.

5 Interestingly, our results are not consistent with those published by Romanazzi et al.<sup>15</sup>, who  
6 found a positive effect of BTH in inducing recovery of BN-infected grapevines. Several hypotheses  
7 can explain these different results: i) the resistance response elicited by BTH is active against BNP  
8 but not against FDP infection ii) different environmental conditions of the two vineyards (in  
9 northern and central Italy, characterized by a very different amount of summer rains) may  
10 influence plant response to the resistance elicitor iii) the grapevine cultivar (Barbera vs  
11 Chardonnay) is a clue element in BTH-elicited defense response. In Sardinia (Italy), the resistance  
12 elicitor Kendal (Valagro, Italy), containing glutathione and oligosaccharines, also failed to promote  
13 recovery of BN-infected Chardonnay and Vermentino grapevines<sup>43</sup>. It is interesting to note that  
14 BTH is active in inducing resistance of grapevines toward other pathogens, such as *Plasmopara*  
15 *viticola* and *Erysiphe necator* (downy and powdery mildews)<sup>44</sup> and *Botrytis cinerea* (gray mold)<sup>45-46</sup>.  
16 These data suggest that BTH induces resistance response in grapevine but this response is  
17 pathogen-specific.

18 Due to the above results, in spite of the great interest in natural, synthetic and biotic  
19 resistance elicitors, a lot of caution should be taken in promoting the application of these tools and  
20 specific evaluations are required for each pathogen-plant combination. Interestingly, in this work  
21 BTH reduced FDP transmission by the vector but was ineffective in inducing recovery of the  
22 infected vines. It is then likely that physiological and molecular mechanisms regulating FDP-  
23 grapevine interactions during early and late stage of plant infection are different, and BTH is only  
24 able to interfere with the former ones. The information on the mechanisms of plant response to

1  
2  
3 1 phytoplasmas are still very limited; only a deeper knowledge of physiological and molecular  
4  
5 2 relationships between the pathogen and the host plant will possibly allow manipulation of such  
6  
7  
8 3 mechanisms, thus providing new effective and environmental friendly tools for the control of  
9  
10 4 phytoplasma diseases of crops.  
11  
12  
13  
14  
15 5  
16 6  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

1

2

3 1 ACKNOWLEDGEMENTS

4

5 2 This work was funded by the Piemonte Region under the grant “Studi sui fattori che favoriscono le

6

7 3 epidemie di flavescenza dorata in Piemonte e loro superamento (FLADO)”. We thank the vine

8

9 4 grower farm F.Ili Bava, Cocconato (Asti, Italy) and the vine grower association Vignaioli Piemontesi

10

11 5 (Castagnito, Italy) for hosting field trials. We also thank dr Ivana Gribaudo, CNR-IPSP, for providing

12

13 6 *in vitro* propagated grapevine plants.

14

15

16

17

18

19

20

21

22 8 REFERENCES

23

24 9 1. Oostendorp M, Kunz W, Dietrich B, Staub T, Induced disease resistance in plants by

25 10 chemicals. *Europ J Plant Pathol* **107**: 19-28 (2001).

26 11 2. Gozzo F, Systemic acquired resistance in crop protection: from nature to a chemical

27 12 approach. *J. Agric Food Chem* **51**: 4487-4503 (2003).

28 13 3. Durrant WE and Dong X, Systemic acquired resistance. *Ann Rev Phytopathol* **42**: 185-209

29 14 (2004).

30 15 4. Stichter L, Mauch-Mani B, and Métraux JP, Systemic acquired resistance. *Ann Rev*

31 16 *Phytopathol* **35**: 235-270 (1997).

32 17 5. Ryals JA ,Neuenschwander UH, Willits MG, Molina A, Steiner HY, Hunt MD, Systemic

33 18 acquired resistance. *Plant Cell* **8**: 1809-1819 (1996).

34 19 6. Pieterse CMJ and van Loon L.C. Signalling cascades involved in induced resistance, in

35 20 *Induced resistance for plant disease control: a sustainable approach to crop protection*, ed.

36 21 by Walters D, Newton A, Lyon G. Blackwell Publishing, Oxford, UK, pp. 65-88 (2007).

37 22 7. Smith JL, De Moraes CM, and Mescher MC, Jasmonate-and salicylate-mediated plant

38 23 defense responses to insect herbivores, pathogens and parasitic plants. *Pest Manag Sci* **65**,

39 24 497-503 (2009).

40 25 8. Pieterse CMJ, van Pelt JA, van Wess SCM, Ton J, Leon-Kloosterziel KM, Keurentjes JJB,

41 26 Verhagen BWM, Knoester M, Van der Sluis L,, Bakker PAHM and van Loon LC, Rhizobacteria-

42 27 mediated induced systemic resistance: triggering, signaling and expression. *Europ J Plant*

43 28 *Pathol* **107**: 51-61 (2001).

44 29 9. Friedrich L, Lawton K, Ruess W, Masner P, Specker N, GutRella M, Meier B, Dincher S, Staub

45 30 T, Uknes S, Metraux JP, Kessmann H, and Ryals J, A benzothiadiazole derivative induces

46 31 systemic acquired resistance in tobacco. *Plant J* **10**:61–70 (1996).

47 32 10. Lawton K, Friedrich L, Hunt M, Weymann K, Delaney T, Kessmann H, Staub T, and Ryals J,

48 33 Benzothiadiazole induces disease resistance in *Arabidopsis* by activation of the systemic

49 34 acquired resistance signal transduction pathway. *Plant J* **10**:71–82 (1996).

50 35 11. Gorlach J, Volrath S, Knauf-Beiter F, Hengy G, Beckhove U, Kogel KH, Oostendorp M, Staub

51 36 T, Ward E, Kessmann H, and Ryals J, Benzothiadiazole, a novel class of inducers of systemic

52 37 acquired resistance, activates gene expression and disease resistance in wheat. *Plant Cell*

53 38 **8**:629–643 (1996).

54 39 12. Romero AM, Kousik CS, and Ritchie DF, Resistance to bacterial spot in bell pepper induced

55 40 by acibenzolar-S-methyl. *Plant Dis* **85**:189–194 (2001).

56

57

58

59

60

13. Bressan A, and Purcell AH, Effect of benzothiadiazole on transmission of X-disease phytoplasma by the vector *Colladonus montanus* to *Arabidopsis thaliana*, a new experimental host plant. *Plant Dis* **89**: 1121-1124 (2005).
14. D'Amelio R, Marzachi C, and Bosco D, Activity of benzothiadiazole on chrysanthemum yellows phytoplasma ('*Candidatus* Phytoplasma asteris') infection in daisy plants. *Crop Prot* **29**:1094-1099 (2010).
15. Romanazzi G, D'Ascenzo D, and Murolo S, Field treatment with resistance inducers for the control of grapevine Bois Noir. *J Plant Pathol* **91**(3): 677-682 (2009).
16. EFSA PLH Panel (EFSA Panel on Plant Health), Scientific Opinion on pest categorisation of Grapevine Flavescence Dorée. *EFSA Journal* **12**(10): 3851, 31 pp. doi:10.2903/j.efsa.2014.3851 (2014).
17. Schvester D, Carle P, Moutous G, Transmission de la flavescence dorée de la vigne par *Scaphoideus littoralis* Ball. *Annales des Epiphyties* **14**: 175-198 (1963).
18. Caudwell A, Kuszala C, Larrue J, and Bachelier JC, Transmission de la Flavescence dorée de la fève à la fève par des cicadelles des genres *Euscelis* et *Euscelidius*. Intervention possible de ces insectes dans l'épidémiologie du Bois noir en Bourgogne. *Ann Phytopathol* hors série: 181-189 (1972).
19. Martini M, Botti S, Marcone C, Marzachi C, Casati P, Bianco PA, Benedetti R, and Bertaccini A, Genetic variability among Flavescence doree phytoplasmas from different origins in Italy and France. *Molec Cell Probes* **16**, 197-208 (2002).
20. Deng S, and Hiruki C, Amplification of 16S rRNA genes from culturable and unculturable molicutes. *Microbiol Methods* **14**: 52-64 (1991).
21. Schneider B, Cousins MT, Klinkong S, and Seemuller E, Taxonomic relatedness and phylogenetic positions of phytoplasma associated with diseases of faba bean, sunnbean, sesame, soya bean and eggplant. *J Plant Dis Prot* **102**: 225-232 (1995).
22. Lee ML, Gundersen DE, Hammond RW and Davis RE, Use of mycoplasma like organism (MLO) group specific oligonucleotide primers for nested-PCR assays to detect mixed MLO infection in a single host- plant. *Phytopathology* **84**:559-566 (1994).
23. Gribaudo I, Cuzzo D and Mannini F, Risanamento da virus e micropropagazione di *Vitis vinifera*. In *Culture in Vitro: Note di Laboratorio*, ed by Micheli M, Caboni E, Lambardi M, Monticelli S, SOI-Società di Ortoflorofrutticoltura Italiana, Sesto Fiorentino (FI), I, 87-89 (2015).
24. Marzachi C, Alma A, D'Aquilio M, Minuto G and Boccardo G, Detection and identification of phytoplasmas infecting cultivated and wild plants in Liguria (Italian Riviera). *J Plant Pathol* **81**: 127-36 (1999).
25. Marzachi C, Veratti F and Bosco D, Direct PCR detection of phytoplasmas in experimentally infected insects. *Ann appl biol* **133**:45-54 (1998).
26. Roggia C, Cagiagli P, Galetto L, Pacifico D, Veratti F, Bosco D and Marzachi C, Flavescence doree phytoplasma titer in field infected Barbera and Nebbiolo grapevines. *Plant Pathol* **63**: 31-41 (2013).
27. Marzachi C, and Bosco D, Relative quantification of chrysanthemum yellows (16Sr I) phytoplasma in its plant and insect host using real-time polymerase chain reaction. *Molec Biotechnol* **30**: 117- 127 (2005)
28. Gamalero E, Marzachi C, Galetto L, Veratti F, Massa N, Bona E, Novello G, Glick BR, Ali S, Cantamessa S, D'Agostino G, Berta G, An 1-aminocyclopropane-1-carboxylate (ACC) deaminase-expressing endophyte increases plant resistance to Flavescence Dorée phytoplasma infection. *Plant Biosystems*, in press. doi: 10.1080/11263504.2016.1174172.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

29. Romanazzi G, Musetti R, Marzachi C, and Casati P, Induction of resistance in the control of phytoplasma diseases. *Petria* **19** (3), 113-129 (2009).

30. Palermo S, Arzone A and Bosco D, Vector-pathogen-host plant relationships of chrysanthemum yellows (CY) phytoplasma and the vector leafhoppers *Macrostelus quadripunctulatus* and *Euscelidius variegatus*. *Entomol exp appl* **99**: 347-354 (2001).

31. Caudwell A, Kuszala C, Bachelier JC, and Larrue J, Transmission de la Flavescence dorée de la vigne aux plantes herbacées par l'allongement du temps d'utilisation de la cicadelle *Scaphoideus littoralis* Ball et l'étude de sa survie sur un grand nombre d'espèces végétales. *Ann Phytopathol* **2**: 415- 428 (1970).

32. Jarausch W, Angelini E, Eveillard S, Malembic-Maher S, Management of fruit tree and grapevine phytoplasma diseases through genetic resistance. In “New Perspectives in Phytoplasma Disease Management”, COST action FA0807 Workshop, Barcelona, Spain, March 22<sup>nd</sup> 2013, pp 56-63 (2013).

33. Carraro L, Ermacora P, Loi N, and Osler R, The recovery phenomenon in apple proliferation-infected trees. *J Plant Pathol* **86**: 141- 146. (2004).

34. Musetti R, Di LS, Ermacora P and Favali MA, Recovery in apple trees infected with the apple proliferation phytoplasma: an ultrastructural and biochemical study. *Phytopathology* **94**: 203–208 (2004).

35. Musetti R., di Toppi LS, Martini M, Ferrini F, Loschi A, and Favali MA, Hydrogen peroxide localization and antioxidant status in the recovery of apricot plants from European Stone Fruit Yellows. *Europ J Plant Pathol* **112**: 53-61 (2005).

36. Seemueller E, Kunze L, and Schaper U, Colonization behaviour of MLO, and symptom expression of proliferation-diseased apple trees and decline-diseased pear trees over a period of several years. *J Plant Dis Prot* **91**: 525–532 (1984).

37. Caudwell A, Les phénomènes de rétablissement chez la Flavescence dorée de la vigne. *Annales des Epiphyties* **12**: 347–354 (1961).

38. Bellomo C, Carraro L, Ermacora P, Pavan F, Osler R, Frausin C and Governatori G, Recovery phenomena in grapevines affected by grapevine yellows in Friuli Venezia Giulia. *Bull Insectol* **60**: 235–236 (2007).

39. Musetti R, Marabottini R, Badiani M, Martini M, di Toppi LS, Borselli S, Borgo M and Osler R, On the role of H<sub>2</sub>O<sub>2</sub> in the recovery of grapevine (*Vitis vinifera* cv. Prosecco) from Flavescence dorée disease. *Functional Plant Biol* **34**(8): 750-758 (2007).

40. Romanazzi G, Murolo S, and Feliziani E, Effects on an innovative Strategy to contain grapevine Bois noir: Field treatment with resistance inducers. *J Phytopathol* **103**: 785-791 (2013).

41. Romanazzi G, and Murolo S, Partial uprooting and pulling to induce recovery in Bois noir infected grapevines. *J Phytopathol* **156**: 747-750 (2008).

42. Schmidt S, Baric S, Massenz M, Letschka T, Vanas V, Wolf M, Kerschbamer C, Zelger R and Schweigkofler W, Resistance inducers and plant growth regulators show only limited and transient effects on infection rates, growth rates and symptom expression of apple trees infected with ‘*Candidatus Phytoplasma mali*’. *J Plant Dis Protect*, 122(5/6), 207-214 (2015).

43. Garau R, Prota VA, Sechi A and Moro G, Biostimulants distribution to plants affected by ‘Bois noir’: results regarding recovery. *Petria* **18**: 366-368 (2008).

44. Dufour MC, Lambert C, Bouscaut J, Mérillon JM, and Corio-Costel MF, Benzothiadiazole-primed defence responses and enhanced differential expression of defence genes in *Vitis vinifera* infected with biotrophic pathogens *Erysiphe necator* and *Plasmopara viticola*. *Plant Pathol* **62**:370- 382 (2013).

- 1  
2  
3 1 45. Iriti M, Rossoni M, Borgo M and Faoro F, Benzothiadiazole enhances resveratrol and  
4 2 anthocyanin biosynthesis in grapevine, meanwhile improving resistance to Botrytis cinerea.  
5 3 *J. Agric Food Chem* **52**: 4406-4413 (2004).  
6 4  
7 46. Iriti M, Rossoni M, Borgo M, Ferrara L and Faoro F, Induction of resistance to gray mold with  
8 5 Benzothiadiazole modifies amino acid profile and increases proanthocyanidins in grape:  
9 6 primary versus secondary metabolism. *J. Agric Food Chem* **53**: 9133- 9139 (2005)  
10  
11 7  
12  
13 8  
14  
15 9  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1:** Number of FDP-infected plants over the total number of grapevines recorded every year among BTH or control treatment plots in the three-year experiment.

	Treatment		
YEAR	BTH	CONTROL	P
2011	4/32	1/26	0.367 (Fisher exact test)
2012	4/41	12/35	0.020 (Chi-square)
2013	0/40	5/40	0.055 (Fisher exact test)

For Peer Review



**Table 2:** Number of FDP-infected plants over the total number of *in vitro* propagated grapevines and estimated amount of FDP cells per ng of vine DNA (mean  $\pm$  s.e.) in BTH-treated and control grapevines at 5 and 11 weeks post inoculation (wpi). Within columns and within rows, different letters indicate significant differences.

Treatment	Number of FD infected plants	FDP cells/ng vine DNA	
		5 wpi	11 wpi
<b>BTH</b>	8/10	3.60 $\pm$ 1.47 <sup>a</sup>	114.56 $\pm$ 56.53 <sup>b</sup>
<b>CONTROL</b>	6/10	1.03 $\pm$ 0.47 <sup>a</sup>	21.43 $\pm$ 7.50 <sup>b</sup>

**Table 3:** Number of recovered plants over the total number of FDP-infected field grapevines at the end of the three-year experiment and estimated amount of FDP cells per ng of vine DNA (mean ± s.e.) in the grapevines still infected at the end of the experiment in BTH-treated and control grapevines.

Treatment	Number of recovered/infected plants	FDP cells/ ng vine DNA
BTH	21/36	731.12 ± 187.18
CONTROL	25/34	1256.73 ± 405.16




Figure 1: Number of FDP-infected grapevines in the three years of the experiment following BTH or water control treatments.

**Supporting Table 1:** FDP-infected or recovered grapevines in the BTH-treatment over the three-year period. Each line represents the sanitary status of an individual plant. +, FDP-infected grapevine; -, recovered grapevine.

2011	2012	2013
+	+	-
+	-	-
+	-	-
+	+	+
+	-	-
+	+	+
+	+	+
+	-	-
+	+	+
+	+	+
+	-	+
+	+	-
+	+	+
+	-	-
+	-	-
+	+	+
+	+	-
+	+	+
+	-	-
+	+	+
+	-	-
+	+	-
+	+	-
+	+	+
+	+	-
+	-	-
+	+	+
+	+	+
+	-	-
+	-	-
+	+	-
+	+	-
+	-	+
+	+	-
+	-	-
+	+	+
+	+	+
+	+	-
+	+	+

**Supporting Table 2:** FDP-infected or recovered grapevines in the control treatment over the three-year period. Each line represents the sanitary status of an individual plant. +, FDP-infected grapevine; -, recovered grapevine.

2011	2012	2013
+	+	-
+	-	-
+	-	-
+	+	-
+	-	-
+	-	-
+	-	-
+	-	-
+	-	-
+	+	-
+	+	+
+	-	-
+	-	-
+	-	-
+	+	-
+	-	-
+	+	+
+	+	+
+	+	+
+	-	-
+	-	+
+	+	+
+	-	+
+	-	-
+	-	-
+	-	-
+	-	-
+	+	-
+	+	+
+	+	-
+	-	-
+	+	+
+	+	-
+	+	-